

TRANSLATOR'S DECLARATION

I, John F. Moloney, Bsc., MIL., CChem., MRSC., translator to Taylor and Meyer of 20 Kingsmead Road, London, SW2 3JD, Great Britain, verify that I know well both the German and the English language, that I have prepared the attached English translation of 12 pages of a German Patent application in the German language with the title:

Verwendung der Acetylaminosäureracemase aus Amycolatopsis orientalis zur Racemisierung von Carbamoylaminosäuren

identified by the code number 000337 AM at the upper left of each page and corresponding to client/matter number of the law firm of

and that the attached English translation of this document is a true and correct translation of the document attached thereto to the best of my knowledge and belief.

I further declare that all statements made of my own knowledge are true and that all statements made on information and belief are believed to be true, and further that these statements are made with the knowledge that wilful false statements and the like are punishable by fine or imprisonment, or both, under 18 USC 1001, and that such false statements may jeopardize the validity of this document.

By: J.F. Moloney

Date: 31st July 2003

Use of acetylamino acid racemase from *Amycolatopsis*
orientalis for racemisation of carbamoylamino acids

The present invention relates to the use of an N-acetyl-
amino acid racemase (AAR) in a process for the racemisation
5 of N-carbamoylamino acids.


Optically pure amino acids are important starting materials
for chemical synthesis and for parenteral nutrition. Many
possibilities of preparing optically pure amino acids are
known to the skilled person. Enzymatic processes, i.a. are
10 suitable in this respect since, on the one hand, they
operate catalytically and on the other hand permit the
preparation of the amino acids with very high enantiomer
enrichment.

A known enzymatic process starts from racemic hydantoins
15 which are transformed to N-carbamoyl-protected amino acids
by means of hydantoinases. These are then converted by
carbamoylases to the amino acids.

The separation of the racemates occurring in this reaction
sequence takes place preferably on the basis of the N-
20 carbamoyl-protected amino acids because both L and D-
selective carbamoylases are available (Park et al.,
Biotechnol. Prog. 2000, 16, 564-570; May et al., Nat
Biotechnol. 2000, 18, 317-20; Pietzsch et al., J.
Chromatogr. B Biomed. Sci. Appl. 2000, 737, 179-86; Chao et
25 al., Biotechnol. Prog. 1999, 15, 603-7; Wilms et al., J.
Biotechnol. 1999, 63, 101-13; Batisse et al., Appl.
Environ. Microbiol. 1997, 63, 763-6; Buson et al., FEMS
Microbiol. Lett. 1996, 145, 55-62).

In order to guarantee complete conversion of the hydantoins
30 used to optically pure amino acids, the necessary
racemisation has taken place hitherto on the basis of
hydantoins by chemical or enzymatic means (EP 745678; EP
542098; scheme 1).

Scheme 1:



N-acetyl amino acid racemases (AARs) from *Streptomyces*
 5 *atratus* Y-53 (Tokuyama et al., Appl. Microbiol. Biotechnol.
 1994, 40, 835-840) and *Amycolatopsis* sp. TS-1-60 (Tokuyama
 et al., Appl. Microbiol. Biotechnol. 1995a, 42, 853-859)
 and *Amycolatopsis orientalis* sp. *lurida* (DE19935268) are
 known. TS-1-60, however, is found to have a very low
 10 activity in the case of N-carbamoyl-protected amino acids.
 Moreover, this enzyme has the disadvantage of a very high
 metal ion dependence, which appears to be a drawback for
 the use of this enzyme in an industrial-scale process.


The object of the present invention was, therefore, to show
 15 the use of an N-acetyl amino acid racemase for the improved
 racemisation of N-carbamoyl amino acids compared with the
 prior art. The intention was that this racemase might be
 used advantageously on an industrial scale in a process for
 the preparation of optically pure amino acid starting from
 20 racemic hydantoins.

The object is achieved by the use of the AAR according to
 claim 1. Claims 2 and 3 relate to preferred embodiments of
 the racemisation process according to the invention.

Due to the fact that an N-acetylamino acid racemase (AAR) from *Amycolatopsis orientalis* subspecies *lurida* (seq. 2) is used in a process for the racemisation of N-carbamoylamino acids, and in view of the surprisingly high activity of the AAR used according to the invention compared with TS-1-60 in terms of the racemisation of N-carbamoylamino acids, it is possible to achieve an equilibrium of enantiomers of N-carbamoyl-protected amino acids in an improved process.

This is particularly advantageous in that it is thus possible to establish a further enzymatic step in a process for the preparation of optically pure amino acids which is based on hydantoins (scheme 2).

Scheme 2:



In contrast to the enzymatic processes known from the literature and which proceed by way of enzymatic or optionally stressing chemical racemisation of hydantoins (scheme 1), a further advantageous possibility of generating optically pure amino acids from racemic hydantoins has thus been created.

The variant of AAR from *Amycolatopsis o. sp. lurida* prepared by recombinant technology according to DE19935268 is preferably used for the racemisation process. It is

known from DE19935268 that this exhibits relatively little heavy metal ion dependence (particularly with regard to cobalt ions) and has low amino acid inhibition. The generation thereof as a recombinant enzyme is also explained therein.

The process according to the invention, as has been mentioned, is used advantageously in an overall process for the preparation of enantiomerically enriched amino acids or derivatives thereof starting from hydantoins or N-carbamoylamino acids. In the case of hydantoins, it is preferable to proceed in such a manner that racemic hydantoins are cleaved by hydantoinases into the corresponding racemic N-carbamoylamino acids and these are then converted by L- or D-specific carbamoylases into the optically active L- or D-amino acids. To ensure that no enrichment of the unconverted enantiomer of an N-carbamoylamino acid takes place in the reaction mixture, the enantiomers of the N-carbamoylamino acids are brought into equilibrium by the addition of the AAR according to the invention and it is thus likewise possible to convert the racemic hydantoin wholly to optically pure amino acids.

This process takes place preferably in an enzyme-membrane reactor (DE 199 10 691.6).

The enzymes mentioned may be used together or successively in the free form as homogeneously purified compounds or as enzymes prepared by recombinant technology. Moreover, the enzymes may also be used as a constituent of a guest organism (whole-cell catalyst as in US09/407062) or in conjunction with the digested cell mass of the host organism. It is also possible to use the enzymes in the immobilised form (Bhavender P. Sharma, Lorraine F. Bailey and Ralph A. Messing, "Immobilisierte Biomaterialien - Techniken und Anwendungen", Angew. Chem. 1982, 94, 836-852). Immobilisation takes place advantageously by freeze-

drying (Dordick et al. J. Am. Chem. Soc. 194, 116, 5009-5010; Okahata et al. Tetrahedron Lett. 1997, 38, 1971-1974; Adlercreutz et al. Biocatalysis 1992, 6, 291-305). Freeze-drying in the presence of surfactant substances such as
5 Aerosol OT or polyvinylpyrrolidone or polyethylene glycol (PEG) or Brij 52 (diethylene glycol monocetyl ether) (Goto et al. Biotechnol. Techniques 1997, 11, 375-378) is more particularly preferred.

The microorganism *Amycolatopsis orientalis* subsp. *lurida* is
10 deposited with the German Collection for Microorganisms under number DSM43134.

The term AAR within the context of the invention means both the native enzyme and the enzyme prepared by recombinant technology.

15 The term enantiomerically enriched denotes the presence of one enantiomer in the mixture with the other in a proportion of >50%.

The term amino acid within the context of the invention means a natural or non-naturally occurring α -amino acid,
20 i.e., the radical situated on the α -C-atom of the α -amino acid may be derived from a natural amino acid as described in Beyer-Walter, Lehrbuch der organischen Chemie, S. Hirzel Verlag Stuttgart, 22nd edition, 1991, p.322f. or also from corresponding α -radicals of non-naturally occurring amino
25 acids which are listed, e.g. in DE19903268.8.

SEQUENCE PROTOCOL

<110> Degussa-Hüls AG

5 <120> Use of an acetylamino acid racemase for the
racemisation of carbamoylamino acids

<130> 000337 AM

10 <140>

<141>

<150> 2

15 <170> PatentIn Ver. 2.1

<210> 1

<211> 1107

<212> DNA

20 <213> Amycolatopsis orientalis

<220>

<221> CDS

<222> (1)..(1107)

25

<400> 1

gtg	aaa	ctc	agc	ggt	gtg	gaa	ctg	cgc	cgg	gtc	cgg	atg	ccg	ctc	gtg	48
Val	Lys	Leu	Ser	Gly	Val	Glu	Leu	Arg	Arg	Val	Arg	Met	Pro	Leu	Val	
1				5					10					15		

30

gcc	ccg	ttc	cgg	acg	tgc	ttc	ggg	acg	cag	tcc	gag	cgg	gaa	ttg	ctg	96
Ala	Pro	Phe	Arg	Thr	Ser	Phe	Gly	Thr	Gln	Ser	Glu	Arg	Glu	Leu	Leu	
			20				25						30			

35

ctg	gtc	cgc	gcg	gtg	acc	ccg	gcg	ggc	gag	ggc	tgg	ggc	gaa	tgt	gtc	144
Leu	Val	Arg	Ala	Val	Thr	Pro	Ala	Gly	Glu	Gly	Trp	Gly	Glu	Cys	Val	
		35					40					45				

40

gcg	atg	gag	gcg	ccg	ctc	tac	tgc	tgc	gag	tac	aac	gac	gcc	gcc	gag	192
Ala	Met	Glu	Ala	Pro	Leu	Tyr	Ser	Ser	Glu	Tyr	Asn	Asp	Ala	Ala	Glu	
	50					55					60					

45

cac	gtg	ctg	cgg	aac	cat	ctg	atc	ccc	gca	ctg	ctg	gcg	gcc	gag	gac	240
His	Val	Leu	Arg	Asn	His	Leu	Ile	Pro	Ala	Leu	Leu	Ala	Ala	Glu	Asp	
	65				70					75				80		

50

gtg	acc	gcg	cac	aag	gtg	acg	ccg	ttg	ctg	gcg	aag	ttc	aag	ggc	cac	288
Val	Thr	Ala	His	Lys	Val	Thr	Pro	Leu	Leu	Ala	Lys	Phe	Lys	Gly	His	
				85				90						95		

cgg	atg	gcg	aag	ggc	gcg	ctg	gag	atg	gcg	gtc	ctc	gac	gcc	gaa	ctc	336
Arg	Met	Ala	Lys	Gly	Ala	Leu	Glu	Met	Ala	Val	Leu	Asp	Ala	Glu	Leu	
		100					105					110				

55

cgc	gcg	cat	gac	cgg	tcc	ttc	gcg	gcc	gag	ctg	ggg	tcc	act	cgc	gac	384
Arg	Ala	His	Asp	Arg	Ser	Phe	Ala	Ala	Glu	Leu	Gly	Ser	Thr	Arg	Asp	
		115					120					125				

tcc	gtg	gcc	tgc	ggg	gtc	tgc	gtc	ggg	atc	atg	gac	tgc	atc	ccg	cac	432
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

	Ser	Val	Ala	Cys	Gly	Val	Ser	Val	Gly	Ile	Met	Asp	Ser	Ile	Pro	His	
	130						135					140					
5	ctg	ctc	gac	gtc	gtc	ggc	ggc	tac	ctc	gac	gag	ggc	tac	gtc	cgg	atc	480
	Leu	Leu	Asp	Val	Val	Gly	Gly	Tyr	Leu	Asp	Glu	Gly	Tyr	Val	Arg	Ile	
	145					150					155					160	
10	aag	ctg	aag	atc	gag	ccc	ggc	tgg	gac	gtc	gag	ccg	gtc	cgg	cag	gtg	528
	Lys	Leu	Lys	Ile	Glu	Pro	Gly	Trp	Asp	Val	Glu	Pro	Val	Arg	Gln	Val	
					165					170					175		
15	cgt	gag	cgc	ttc	ggc	gac	gac	gtg	ctg	ctg	cag	gtc	gac	gcg	aac	acc	576
	Arg	Glu	Arg	Phe	Gly	Asp	Asp	Val	Leu	Leu	Gln	Val	Asp	Ala	Asn	Thr	
					180					185					190		
20	gcg	tac	acg	ctg	ggc	gac	gcg	ccc	ctg	ctg	tcc	cgg	ctc	gac	ccg	ttc	624
	Ala	Tyr	Thr	Leu	Gly	Asp	Ala	Pro	Leu	Leu	Ser	Arg	Leu	Asp	Pro	Phe	
			195					200					205				
25	gac	ctg	ctg	ctg	atc	gag	cag	ccg	ctc	gaa	gaa	gag	gac	gtg	ctc	ggc	672
	Asp	Leu	Leu	Leu	Ile	Glu	Gln	Pro	Leu	Glu	Glu	Glu	Asp	Val	Leu	Gly	
		210					215					220					
30	cac	gcc	gag	ctg	gcc	aag	cgg	atc	cgg	acg	ccg	atc	tgc	ctc	gac	gag	720
	His	Ala	Glu	Leu	Ala	Lys	Arg	Ile	Arg	Thr	Pro	Ile	Cys	Leu	Asp	Glu	
	225					230					235					240	
35	tgc	atc	gtc	tgc	gcc	aag	gcc	gcc	gcg	gac	gcg	atc	aag	ctc	ggc	gcc	768
	Ser	Ile	Val	Ser	Ala	Lys	Ala	Ala	Ala	Asp	Ala	Ile	Lys	Leu	Gly	Ala	
					245					250					255		
40	tgc	cag	atc	gtc	aac	atc	aaa	ccg	ggc	cgg	gtc	ggc	gga	tac	ctc	gaa	816
	Cys	Gln	Ile	Val	Asn	Ile	Lys	Pro	Gly	Arg	Val	Gly	Gly	Tyr	Leu	Glu	
				260					265					270			
45	gcc	cgc	cgg	gtg	cac	gac	gtc	tgc	gcg	gca	cac	ggg	atc	gcg	gtg	tgg	864
	Ala	Arg	Arg	Val	His	Asp	Val	Cys	Ala	Ala	His	Gly	Ile	Ala	Val	Trp	
			275					280					285				
50	tgc	ggc	ggg	atg	atc	gag	acc	ggg	ctc	ggc	cgg	gcg	gcc	aac	gtc	gca	912
	Cys	Gly	Gly	Met	Ile	Glu	Thr	Gly	Leu	Gly	Arg	Ala	Ala	Asn	Val	Ala	
		290					295					300					
55	ctg	gcc	tgc	ctg	ccc	ggc	ttc	acg	ctg	ccg	ggg	gac	acc	tgc	gcg	tcc	960
	Leu	Ala	Ser	Leu	Pro	Gly	Phe	Thr	Leu	Pro	Gly	Asp	Thr	Ser	Ala	Ser	
	305					310					315					320	
60	ggc	cgg	ttc	tat	cgc	acc	gac	atc	acc	gag	ccg	ttc	gtg	ctg	gac	gcc	1008
	Gly	Arg	Phe	Tyr	Arg	Thr	Asp	Ile	Thr	Glu	Pro	Phe	Val	Leu	Asp	Ala	
					325					330					335		
65	ggg	cat	ctg	ccg	gtg	ccg	acc	ggg	ccg	ggc	ctc	ggg	gtg	act	ccg	att	1056
	Gly	His	Leu	Pro	Val	Pro	Thr	Gly	Pro	Gly	Leu	Gly	Val	Thr	Pro	Ile	
				340					345					350			
70	ccg	gat	ctt	ctg	gac	gag	gtc	acc	acg	gag	aaa	gcg	tgg	atc	ggc	tgc	1104
	Pro	Asp	Leu	Leu	Asp	Glu	Val	Thr	Thr	Glu	Lys	Ala	Trp	Ile	Gly	Ser	
			355					360						365			

tag

1107

5 <210> 2
 <211> 368
 <212> PRT
 <213> Amycolatopsis orientalis

10 <400> 2
 Val Lys Leu Ser Gly Val Glu Leu Arg Arg Val Arg Met Pro Leu Val
 1 5 10 15
 Ala Pro Phe Arg Thr Ser Phe Gly Thr Gln Ser Glu Arg Glu Leu Leu
 20 25 30
 15 Leu Val Arg Ala Val Thr Pro Ala Gly Glu Gly Trp Gly Glu Cys Val
 35 40 45
 Ala Met Glu Ala Pro Leu Tyr Ser Ser Glu Tyr Asn Asp Ala Ala Glu
 50 55 60
 His Val Leu Arg Asn His Leu Ile Pro Ala Leu Leu Ala Ala Glu Asp
 65 70 75 80
 20 Val Thr Ala His Lys Val Thr Pro Leu Leu Ala Lys Phe Lys Gly His
 85 90 95
 Arg Met Ala Lys Gly Ala Leu Glu Met Ala Val Leu Asp Ala Glu Leu
 100 105 110
 25 Arg Ala His Asp Arg Ser Phe Ala Ala Glu Leu Gly Ser Thr Arg Asp
 115 120 125
 Ser Val Ala Cys Gly Val Ser Val Gly Ile Met Asp Ser Ile Pro His
 130 135 140
 30 Leu Leu Asp Val Val Gly Tyr Leu Asp Glu Gly Tyr Val Arg Ile
 145 150 155 160
 Lys Leu Lys Ile Glu Pro Gly Trp Asp Val Glu Pro Val Arg Gln Val
 165 170 175
 Arg Glu Arg Phe Gly Asp Asp Val Leu Leu Gln Val Asp Ala Asn Thr
 180 185 190
 35 Ala Tyr Thr Leu Gly Asp Ala Pro Leu Leu Ser Arg Leu Asp Pro Phe
 195 200 205
 Asp Leu Leu Leu Ile Glu Gln Pro Leu Glu Glu Glu Asp Val Leu Gly
 210 215 220
 His Ala Glu Leu Ala Lys Arg Ile Arg Thr Pro Ile Cys Leu Asp Glu
 225 230 235 240
 40 Ser Ile Val Ser Ala Lys Ala Ala Ala Asp Ala Ile Lys Leu Gly Ala
 245 250 255
 Cys Gln Ile Val Asn Ile Lys Pro Gly Arg Val Gly Gly Tyr Leu Glu
 260 265 270
 45 Ala Arg Arg Val His Asp Val Cys Ala Ala His Gly Ile Ala Val Trp
 275 280 285
 Cys Gly Gly Met Ile Glu Thr Gly Leu Gly Arg Ala Ala Asn Val Ala
 290 295 300
 50 Leu Ala Ser Leu Pro Gly Phe Thr Leu Pro Gly Asp Thr Ser Ala Ser
 305 310 315 320
 Gly Arg Phe Tyr Arg Thr Asp Ile Thr Glu Pro Phe Val Leu Asp Ala
 325 330 335
 Gly His Leu Pro Val Pro Thr Gly Pro Gly Leu Gly Val Thr Pro Ile
 340 345 350
 55 Pro Asp Leu Leu Asp Glu Val Thr Thr Glu Lys Ala Trp Ile Gly Ser
 355 360 365

Examples:

Detection of racemase activity of the recombinant AAR enzyme

The substrate spectrum of the N-acetylamino acid racemase
5 from *Amycolatopsis orientalis* subsp. *lurida* was tested using the enzyme assay described below.

The assay was composed of the following:

Tris/HCl buffer	50 mM (pH 8.0)
10 Substrate	25 mM
Cobalt chloride	6 mM
AAR	approx 150 μ g purified protein
Final volume	1 ml

Enantiomerically pure amino acid derivatives were used in
15 the test and the formation of the corresponding racemate was monitored in the polarimeter (Perkin-Elmer 241). Incubation took place at 30°C (heated cell) for 3 to 12 hours. The measurements were taken at a wavelength λ = 365 nm.

Table 1: List of the substrates tested and of the corresponding specific activity of the AAR.

Substrate	Specific activity
N-Carbamoyl-D-Met	155 mU/mg
N-Carbamoyl-D-Phe	20 mU/mg
N-Carbamoyl-L-Abs	15 mU/mg
N-Carbamoyl-L-Leu	20 mU/mg
N-Carbamoyl-L-Met	118 mU/mg
N-Carbamoyl-L-Tyr	62 mU/mg
N-Carbamoyl-L-Val	20 mU/mg

5

The N-acyl amino acid racemase from A. TS-1-60 with N-carbamoyl-D-Met as substrate has an activity of 100 mU/mg. This specific activity is thus 35% lower than that of the racemase from A. orientalis subsp. lurida.

10

Patent claims:

1. Use of N-acetylamino acid racemases (AAR) from
Amycolatopsis orientalis subspecies lurida in a
process for the racemisation of N-carbamoylamino
5 acids.
2. The use as claimed in claim 1 in a process for the
preparation of enantiomerically enriched amino acids
or derivatives thereof starting from hydantoins or N-
carbamoylamino acids.
- 10 3. The use as claimed in one of the preceding claims,
wherein
the process is carried out in an enzyme-membrane
reactor.

Abstract:

The invention relates to the use of the N-acetylamino acid racemase from *Amycolatopsis orientalis* subspecies *lurida* for the racemisation of N-carbamoylamino acids.

- 5 This use permits the 100% preparation of optically pure amino acids starting from racemic hydantoins in an enzymatic overall process.